

Stratified simplices and intersection homology

Jonathan Fine*

23 July 1998

Abstract

Intersection homology is obtained from ordinary homology by imposing conditions on how the embedded simplices meet the strata of a space X . In this way, for the middle perversity, properties such as strong Lefschetz are preserved. This paper defines local-global intersection homology groups, that record global information about the singularities of X . They differ from intersection homology in that stratified rather than ordinary simplices are used. An example of such is $\sigma_j \times C\sigma_i$, where σ_i and σ_j are ordinary simplices, and C is the coning operator. The paper concludes with a sketch of the relationship between local-global homology and the geometry of convex polytopes. This paper is a more formal exposition of part of the author's *Local-global intersection homology*, alg-geom/9709011.

1 Introduction

Throughout X will be an irreducible complex algebraic variety of complex dimension n , considered as a topological space of real dimension $m = 2n$. Homology theory, as developed by Poincaré, associates to X a family $H_i X$ of ordinary homology groups, where the index i runs from 0 to m . Throughout, we will calculate homology with real coefficients. When X is nonsingular, as is well known, these homology groups have the following properties.

First, there is defined a natural intersection pairing

$$H_i X \otimes H_j X \rightarrow H_0 X \cong \mathbf{R}$$

whenever $i + j = m$, and this pairing is perfect (non-degenerate). This is *Poincaré duality*. Second, each embedding $X \subset \mathbf{P}_N$ of X in a projective space gives rise to a hyperplane class or *Lefschetz element* $\omega \in H_{m-2} X$ with the following property. There defined is a natural intersection map

$$\omega : H_i X \rightarrow H_{i-2} X$$

that is injective for $i \geq n$ and surjective for $i - 2 \leq n$. Equivalently, for $i + j = m$, $i \geq j$ and $2k = i - j$ the map

$$\omega^k : H_i X \rightarrow H_j X$$

is an isomorphism. This is the *strong Lefschetz theorem*. Thirdly, as a consequence of Deligne's proof of the Weil conjectures, the *Betti numbers* $h_i X$ (the dimension of $H_i X$) can be computed by counting points on X over finite fields, and thus there are combinatorial formulae for the Betti numbers $h_i X$.

*203 Coldhams Lane, Cambridge, CB1 3HY, England. E-mail: j.fine@pmms.cam.ac.uk
This version: 17 September 1998

Now suppose that the algebraic variety X is singular. In this case, as a topological space X need not be a manifold, and in general the properties listed do not hold for the ordinary homology of X . They are true however for the middle perversity intersection homology (mpih) theory introduced by Goresky and MacPherson. This theory is in its definition similar to ordinary homology, except that it imposes restrictions on how the cycles and boundaries meet the singularities. Poincaré duality was proved by Goresky and MacPherson [9]. Strong Lefschetz was proved in [2], again as a consequence of the proof of the Weil conjectures. For toric varieties, see any of [3, 4, 12] for the Betti number formula. Throughout, the middle perversity will be used.

The ordinary homology of nonsingular varieties has other properties, such as the Riemann-Hodge inequalities and a ring structure, whose analogues for the intersection homology of singular varieties is at present unknown. For the author, combinatorial formulae for the Betti numbers are the most significant property, whereas classically it is the functorial nature that is pre-eminent.

2 Ordinary and intersection homology

In this section and the next we review matters, so as to provide some background and motivation.

As is well known, the ordinary homology of X can be defined in the following way. The *standard i -simplex* σ_i is the convex hull of the standard basis vectors $\mathbf{e}_0, \dots, \mathbf{e}_i$ in $\mathbf{R}^{i+1}$. Alternatively, it is the region defined by the inequalities $x_0 \geq 0, \dots, x_i \geq 0$ and the equation $x_0 + \dots + x_i = 1$. An *embedded i -simplex* f is a continuous map $f: \sigma \rightarrow X$.

Each embedded i -simplex f has a *boundary* df , which is a formal sum of embedded $(i-1)$ -simplices. More exactly

$$df = \partial_0 f - \partial_1 f + \dots + (-1)^i \partial_i f$$

where $\partial_j f$ is the composition of f with the j -th *inclusion map* $\sigma_{i-1} \hookrightarrow \sigma_i$. (Each point in σ_{i-1} is a sequence of i numbers. This map prolongs the sequence by inserting a zero immediately before the entry indexed by j , or after the last entry if $j = i$.) It is easily seen that $d^2 f = 0$ for any embedded simplex f . Later, we will need to know what the boundary of a product of ordinary simplices is.

To continue, an *i -chain* ξ is simply a formal sum of embedded i -simplices. A chain η is said to be a *cycle* if its boundary $d\eta$, also as a formal sum, is zero. Because $d \circ d$ is zero, the boundary $d\xi$ of a chain ξ will always be a cycle. A cycle η is called a *boundary* if the equation $\eta = d\xi$ can be solved, for some chain ξ . The i -th *ordinary homology group* $H_i X$ of X is defined to be the quotient group of the i -cycles modulo the i -cycles that are boundaries.

Intersection homology is obtained by imposing conditions on the embedded simplices. To do this, we must first stratify X . Throughout we will suppose that X has been written as a disjoint union

$$X = S_0 \sqcup S_1 \sqcup \dots \sqcup S_n$$

of strata S_j . At each of its points s_j each S_j is to have complex dimension j , and about each s_j the topological structure of X is to be locally constant. The *middle perversity conditions* on an embedded i -simplex f are this. For each $j < n$ the dimension of $f^{-1}(S_j)$ must be at most $(i-1) - (n-j)$. Put another way, there is of course no condition on $f^{-1}(S_n)$, but as j goes from n to 0, so the largest allowed dimension drops first by two, and then by one at each step.

An *allowed* or *admitted i -cycle* η is as before a formal sum of i -chains whose boundary $d\eta$ is zero, where in addition each embedded simplex in η satisfies the above conditions. Such a cycle is an *allowed boundary* if the equation $\eta = d\xi$ can be solved, where the embedded simplices of ξ satisfy the above conditions. The i -th (middle perversity) *intersection homology group* $IH_i X$ of X consists of the allowed i -cycles modulo the allowed boundaries.

3 Local intersection homology

This section continues the review of background material, and concludes with some examples of local-global homology groups.

As mentioned earlier, the (middle perversity) intersection homology groups have the Poincaré duality, strong Lefschetz and combinatorial formula properties. However, when X is singular there can be significant information about the singularities that is not recorded by these groups. This information is recorded after a fashion by intersection homology with other perversities, or by the various ‘change of perversity’ groups that can be defined. But these groups generally fail to satisfy strong Lefschetz, and do not have a combinatorial formulae for the Betti numbers [11]. The local-global theory is an attempt to define additional homology groups, whose Betti numbers are we hope given by a combinatorial formula.

To motivate the definition, we will study the topology of X about a stratum S_j . First let s be a point on S_j . About s the space X looks like the cone on something, and that something is the *link* L_s of X about s . Up to homeomorphism, L_s is locally constant on S_j . Now suppose that η is an i -cycle on L_s , with $i \geq 1$. The space X is about s homeomorphic to CL_s , and so in X the cycle η can be ‘coned away to the apex’. In other words, it is the boundary of the cone $C\eta$ of η , or $\eta = dC\eta$. For ordinary homology this expresses η as a boundary, but for intersection homology the $(i+1)$ -chain $C\eta$ is not allowed. Put another way, ordinary homology is locally trivial, but intersection homology is not.

In this way one can define the local intersection homology groups of X at s . When X is a Schubert variety they have been much studied, and their Betti numbers are given by the celebrated Kazhdan-Lusztig polynomials. These groups do not however record global information about X . To take a step in this direction, we will allow the point s to move. To fix ideas, suppose that s_t , $0 \leq t \leq 1$ is a one dimensional path γ on X . So long as s_t stays on S_j , local constancy allows a local cycle η_0 at s_0 to be moved to another local cycle η_1 at s_1 . Now suppose that s_0 and s_1 are the same point. Although η_0 and η_1 are equivalent along γ , they need not be equivalent in a neighborhood of s_0 . This is an example of the *monodromy action of the fundamental group of S_j on the local homology at $s \in S_j$* .

Now suppose that s_t , $0 \leq t \leq 1$ is again a path γ on X , but this time we will allow s_t to change strata. More exactly, we shall suppose that for $t > 0$ one has $s_t \in S_j$, but that s_0 lies on some other strata S_k . (Necessarily, k will be smaller than j .) Now let η_1 be a local cycle at s_1 . We can again move η_1 along γ to give local cycles η_t at s_t , provided t is not zero. When t is close to zero, η_t will lie in a small ball about s_0 , and so can be thought of as a cycle local to s_0 . Thus, paths from s_j to s_k will take local cycles from S_j to S_k .

Now form the group consisting of all formal sums of local i -cycles on X , modulo the relations due to (a) local boundaries, (b) equivalence along paths lying entirely on strata, and (c) equivalence due to paths as in the previous paragraph. This is a gluing together of local homology groups, and is a global invariant of X . We will show that it does not depend on the choice of a stratification.

First, choose a point $s_{i,j}$ on each connected component of each stratum S_i on X . We are ‘gluing together’ the local homology groups at the $s_{i,j}$. Now refine the stratification. (If need be, first move the $s_{i,j}$ so that they avoid the new components.) This will add new $s_{i,j}$, and may potentially reduce the relations that are available. But each new $s_{i,j}$ will lie on an old stratum connected component, and so the local homology at such a new point will be equivalent to the local homology on the old component.

Now for the relations. Suppose that γ is a path, as previously considered. Just as removing the origin does not disconnect $\mathbf{C}$, so removing subvarieties from X does not destroy the equivalence induced by γ , provided its end points are left intact. This proves the result.

The above definition looks at equivalence of (formal sums of) local cycles over the whole of X . If instead of X one applies the definition to $S_k \cup \dots \cup S_n$, one obtains additional invariants of X . The same arguments as before demonstrates that these groups are stratification independence.

These groups are the simplest significant examples of local-global intersection homology. In the next section we introduce stratified simplices, which are the basic building blocks for local-global cycles, in their topological form. The following section will then define the local-global groups, first without regard to the avoidance of strata, and then with such regard.

4 Stratified simplices

Just as ordinary intersection homology can be defined using embeddings of ordinary simplices σ_i , so local-global homology can be defined using stratified simplices. These objects can also be used to give a formal expression to some of the concepts of the previous section.

For example, suppose that η_1 is an (intersection homology) i -cycle local to the point s of X , with $i \geq 1$. By identifying a ball about s with the cone CL_s on the link L_s at s , one can produce from η_1 a family of η_t of local cycles, for $1 \geq t > 0$, such that in the limit $t = 0$ the ‘cycle’ η_0 is supported by s . This is, as mentioned earlier, an example of the ‘coning away’ of a cycle. Such are allowed by ordinary homology, but are in general inadmissible for intersection homology.

We will now describe this example, without explicit reference to the base point s . First let $f_1 : \sigma_i \rightarrow X$ be one of the embedded simplices out of which the cycle η_1 is constructed. Because η is local (to s) we can extend f_1 to the cone $C\sigma_i$ on the simplex σ_i , to obtain a continuous map $f : C\sigma_i \rightarrow X$. This is an example of an embedded stratified simplex. Its boundary will be defined so that the formal sum corresponding to η_1 will again be a cycle. In other words, the boundary df will be a formal sum of maps from $C\sigma_{i-1}$. Put another way, f is a continuous function $f(t, \lambda)$ of t and λ , where $\lambda \in \sigma_i$ and $t \in [0, 1]$, subject to the constraint that $f(0, \lambda)$ is independent of λ . We take the boundary in the λ directions, but not in the t direction.

Now suppose that η is a formal sum of embedded $C\sigma_i$, such that for each $1 \geq t > 0$ the resulting η_t is an intersection homology i -cycle. This is equivalent to saying that $d\eta$ as just described is zero, and also that each $f(t, \lambda)$ in η is admissible as an embedded i -simplex, for $0 < t \leq 1$. Because $C\sigma_i$ is a cone it follows that for each embedding of $C\sigma_i$ the point $s = f(0, \lambda)$ will be independent of the choice of λ in σ_i . In this way the existence of ‘base points’ is, as promised, implicit in the construction. It should now be clear that if η_1 is a formal sum of local cycles, as used in the previous section, then it can be presented as the $t = 1$ surface of a formal sum η of embedded $C\sigma_i$, and vice versa.

The relations used between these η are more subtle. If η_1 and η'_1 are equivalent as local cycles then there will, by definition, be a formal sum of embedded σ_{i+1} , whose boundary is $\eta_1 - \eta'_1$. This can then be coned to give a formal sum of $C\sigma_{i+1}$, whose boundary is $\eta - \eta'$. Although such relations are perfectly valid, they do not allow the base points to move. (In the previous section locally constancy was used to do this.)

To move η_1 at s to an equivalent η'_1 at s' , we need a path γ from s to s' . This path is a map from $[0, 1]$ to X . Thus, to produce a chain whose boundary is $\eta - \eta'$ it is natural to consider embeddings of $\sigma_1 \times C\sigma_i$, where σ_1 is of course a 1-simplex. If f is an embedding of $\sigma_1 \times C\sigma_i$ then its boundary will be a formal sum of embeddings of $\sigma_0 \times C\sigma_i \cong C\sigma_i$ and also of $\sigma_1 \times C\sigma_{i-1}$. As before, we say that f is *allowed* if for each $0 < t \leq 1$ the embedding f_t of $\sigma_1 \times \sigma_i$ is allowed by the perversity conditions. We will now say that a formal sum η of embedded $C\sigma_i$ is a *boundary* if first it is a chain, and second the equation $\eta = d\xi$ can be solved, where ξ is a formal sum of allowed embedded $C\sigma_{i+1}$ and $\sigma_1 \times C\sigma_i$ simplices. That the stratified simplices used to construct ξ are not all of the

same type is a subtlety not previously present.

The basic operations in the above examples are first applying the cone operator C to a simplex, and second multiplying by an ordinary simplex. We can now give the main definition of this section.

Definition 1 *An order zero stratified simplex is just an ordinary simplex σ . If σ is an order $(r - 1)$ stratified simplex and σ_i is an ordinary simplex (of dimension i) then $\sigma_i \times C\sigma$ is an order r stratified simplex, and all such arise in this way, for $r \geq 1$.*

The dimension sequence (or dimension for short) of an ordinary i -simplex is (i) . The dimension of $\sigma_i \times C\sigma$ is $(\dots, i)$, where the dots denote the entries of the dimension sequence of σ . We will use $\sigma(i_0, \dots, i_r)$ (or more briefly $\sigma(i)$ or even just σ_i) to denote the standard stratified simplex of dimension $i = (i_0, \dots, i_r)$.

5 Local-global intersection homology

The usual concepts, when applied to stratified simplices, will produce some but not all of the local-global intersection homology groups. The remainder arise through the imposition of additional conditions, as to where the ‘apex locii’ of the embedded simplices meet the strata of X .

Throughout this section σ will be an order r stratified simplex of dimension $i = (i_1, \dots, i_r)$. As usual, an *embedding* of σ is just a continuous map $f : \sigma \rightarrow X$. We will say that f is *allowed* if, whenever the coning variables t are all nonzero, the restriction

$$f_t : \sigma(i_r) \times \dots \times \sigma(i_0) \rightarrow X$$

of f to the t -slice is allowed by the perversity conditions.

The *boundary* df of f is defined so that the t -slice $(df)_t$ of df is equal to the boundary $d(f_t)$ of the t -slice of f . This condition is sufficient to determine df . It also ensures that $d \circ d$ is zero. Put another way, the construction is this. The product of simplices $\sigma(i_r) \times \dots \times \sigma(i_0)$ has $i_0 + \dots + i_r + r$ boundary facets, each one of which when suitably coned up produces a facet of σ . (This assumes that none of the i_j are zero. But such can be ignored.) Each facet of a $\sigma(i_j)$ determines a facet of σ . The sign associated to such a facet is to be determined by its location (even or odd) in $\sigma(i_r) \times \dots \times \sigma(i_0)$, and not by its location in $\sigma(i_j)$ itself. This ensures in the usual way that $d \circ d$ is zero. The *boundary* df of $f : \sigma \rightarrow X$ is the signed formal sum of the restriction of f to these boundary facets.

As usual, an *i -cycle* is a formal sum η of allowed embedded i -simplices, whose boundary $d\eta$ is zero. Such a cycle is a *boundary* if the equation $\eta = d\xi$ has a solution, where ξ is a formal sum of allowed embedded stratified simplices. Notice that in this situation it makes no sense to talk of $(i + 1)$ simplices, for i is not an integer.

Definition 2 *Let $i = (i_0, \dots, i_r)$ be a sequence of non-negative integers. The (unrestricted) local-global homology group $H_{i;0}X$ is defined to be the i -cycles modulo the i -boundaries, as defined in the previous paragraph.*

Just as a cone has an apex, so an order r stratified simplex will have r apex locii. These induce a stratification of σ . Further local-global groups can be obtained by imposing conditions on how the strata of σ meet the strata of X . The following definitions make these concepts clear.

Definition 3 *Suppose $\sigma = \sigma_i \times C\sigma'$ is an order r stratified simplex. The r -th apex locus $A_r\sigma$ of σ consists of $\sigma_i \times \{0\}$, where $\{0\}$ is the apex of $C\sigma'$. The j -th apex locus $A_j\sigma$, for $r > j \geq 1$, consists of $\sigma_i \times C(A_j(\sigma'))$, or in other words $\sigma_i \times C$ applied to the j -th apex locus of σ' . We define $A_0\sigma$ to*

be σ itself, but do not count it as an apex locus. The $A_j\sigma$ provide a descending filtration of σ . The successive complements $A_j\sigma \setminus A_{j+1}\sigma$ are called the strata $A_j^o\sigma$ of σ .

Definition 4 Suppose $f : \sigma \rightarrow X$ is an embedded order r stratified simplex. The where or w -sequence $w(f) = (w_1, \dots, w_r)$ of f is defined in the following way. The entry w_i is defined to be the largest j such that $f^{-1}(S_j \cup \dots \cup S_n)$ meets the apex locus $A_i\sigma$ in a set that is dense in $A_i\sigma$.

Definition 5 Let $w = (w_1, \dots, w_r)$ be a w -sequence. Say that an embedding $f : \sigma \rightarrow X$ of an order r simplex is w -allowed if for $1 \leq j \leq r$ the inequality $w_j \leq w_j(f)$ holds.

Definition 6 Let i be an index for order r stratified simplices, and let $w = (w_1, \dots, w_r)$ be a w -sequence. The (i, w) -th local-global intersection homology group $H_{i;w}X$ is the quotient group of i -cycles η modulo i -boundaries $\psi = d\xi$, where each of η , ψ and ξ uses only w -allowed stratified simplices. One can also call these groups, for $r > 0$, the higher order intersection homology groups.

At the end of §4 we proved the following result.

Theorem 7 If i is of the form $(i_1, 0)$, and w is (w_1) , then the (i, w) local-global homology $H_{i;w}X$ of X is independent of the stratification of X .

6 Betti numbers and convex polytopes

One of the most important properties of middle perversity intersection homology is that in many cases there is a combinatorial formula for the Betti numbers. Here we will say a few words about this, and its relation to local-global homology. The basic idea is that from the Betti numbers for a small class of examples, we can reconstruct the homology theory.

First, we need some algebraic varieties. If X is an algebraic variety we will use IX to denote the product of X with $\mathbf{P}_1$, while CX will denote the projective cone on X . (This assumes that X is given to us as a projective variety.) Now let W be a word of length n in I and C , terminated by a $\{\text{pt}\}$. Thus W denotes an n -dimensional projective variety. These IC varieties are in fact toric varieties, of rather a special type.

The mpiv Betti numbers, also known as the h -vector, of IW can be computed from that of W by the Künneth formula. In other words

$$h(IW) = IhW \tag{1}$$

where the I on the right hand side represents convolution with $(1, 1)$, the Betti numbers of $\mathbf{P}_1$. (Because the odd Betti numbers are here all zero, it is convenient in this section to omit them.)

Now consider the cone CW on W . It follows easily from the standard formula for toric variety Betti numbers that $h(CW)$ is obtained from hW by repeating the middle term. In other words, if $C(hW)$ denotes hCW then

$$\begin{aligned} C(a) &= (a, a) \\ C(a, a) &= (a, a, a) \\ C(a, b, a) &= (a, b, b, a) \\ C(a, b, b, a) &= (a, b, b, b, a) \\ C(a, b, c, b, a) &= (a, b, c, c, b, a) \end{aligned} \tag{2}$$

and so on. It is easy to guess that this should be the required formula, although proving its truth is another matter.

We can now apply results in combinatorics. Let Δ be a convex polytope. It has a flag vector $f\Delta$ [1, 5]. If $h\Delta$, the Betti numbers of the associated toric variety $\mathbf{P}_\Delta$, is a linear function of $f\Delta$, then it is determined by its value on the IC polytopes. This holds for any linear function of the flag vector, and not just the $mpih$ h -vector. (Here, $I\Delta$ is the product of Δ with an interval, a *cylinder* or *prism*; while $C\Delta$ is the *cone* or *pyramid* on Δ . The result of applying a word in I and C to the point polytope is an IC polytope.) Thus, the formulae (1) and (2) determine $h\Delta$ for all Δ , provided we either know or assume that $h\Delta$ is indeed a linear function of $f\Delta$.

Now pretend that we know nothing about intersection homology, except that there may be a topological homology theory whose Betti numbers are given by (1) and (2). With a certain amount of work it is possible to produce an explicit recursive formula for $h\Delta$ in terms of $f\Delta$, and when this is done each flag δ on Δ makes a numerical contribution λ_δ to $h\Delta$. These numerical contributions can be interpreted in terms of the linear algebra associated to δ [7], and then in terms of the topology of $\mathbf{P}_\Delta$. The conclusion of this process is that, at least conjecturally, the formulae (1) and (2) give rise to $mpih$ cycles and boundaries on $\mathbf{P}_\Delta$, and hence by way of example tell us what the $mpih$ conditions on embedded simplices are.

Put in a nutshell, the argument is that (1) and (2), together with a certain facility in the combinatorics of Δ and the topology of $\mathbf{P}_\Delta$, would have allowed us to discover middle perversity intersection homology, if only we had thought to follow this path before Goresky and MacPherson discovered it via topology.

In this paper we have given a topological definition of local-global intersection homology. The actual process of discovery was as described in this section. First, rules analogous the the rules (1) and (2) for I and C were hypothesized, and then the definitions were unwound to give the topological definition.

The first step was not easy. Not every pair I and C of rules will give rise to a linear function on the flag vector. In fact, they must satisfy the IC -equation [5]

$$I(I - C)C = (I - C)CI$$

and also the boundary condition $I\{\text{pt}\} = C\{\text{pt}\}$, and these conditions are sufficient.

The reader might wish, as an exercise, to compute perhaps only heuristically the local-global Betti numbers for IC varieties, perhaps only in smallish dimensions. When doing this it is very helpful to know, as a consequence of a fundamental result of Bayer and Billera [1], that one wishes for projective toric varieties of dimension n to have F_{n+1} , the $(n+1)$ -st Fibonacci number, linearly independent Betti numbers. The author's answer to this exercise is the starting point for [6], which contains a fuller and less formal, exposition of concepts presented in this paper.

Acknowledgements

The main definition of this paper is the outcome of a long search that was prompted by the publication in 1985 of [1]. Of the many people who showed a tolerant interest in the author's previous attempts, special thanks are due to Marge Bayer, Gil Kalai, Carl Lee, Peter McMullen and Richard Stanley. The term 'stratified simplex' is due to an anonymous referee of [7].

References

- [1] M.M. Bayer and L.J. Billera, Generalized Dehn-Sommerville equations for polytopes, spheres, and Eulerian ordered sets, *Invent. Math.* **79** (1985), 143–157
- [2] A. Belinson, J. Bernstein and P. Deligne, *Faisceaux pervers*, Asterisque 100, Société mathématique de France, (1982)
- [3] J. Denef and F. Loeser, Weights of exponential sums, intersection homology and Newton polyhedra, *Invent. Math.* **106** (1991), 275–294
- [4] K. Fieseler, Rational intersection homology of projective toric varieties, *J. reine angew. Math.* **413** (1991), 88–98
- [5] J. Fine, The Mayer-Vietoris and *IC* Equations for Convex Polytopes, *Discrete Comput. Geom.* **13** (1995), 177–188
- [6] ———, Local-global intersection homology, preprint alg-geom/9709011 (September 1997)
- [7] ———, Convex polytopes and linear algebra, preprint alg-geom/9710001 (October 1997)
- [8] ———, Ring structure, uniform expressions and intersection homology, preprint math.AG/9805100 (May 1998)
- [9] M. Goresky and R. MacPherson, Intersection homology theory, *Topology* **19** (1980), 135–162
- [10] ———, Intersection homology II, *Invent. Math.* **72** (1983), 77–129
- [11] M. McConnell, The rational homology of toric varieties is not a combinatorial invariant, *Proc. Amer. Math. Soc.* **105** (1989), 986–991
- [12] ———, Generalized *h*-vectors, intersection cohomology of toric varieties and related results, *Adv. Stud. Pure Math.* **11** (1987), 187–213